

## VII Leopoldo García-Colín Mexican Meeting on Mathematical and Experimental Physics



Contribution ID: 67

Type: **not specified**

# HORACIO PASTAWSKI: Decoherent Quantum Transport: from Giant Magnetoresistance and SASERs to Quantum Dynamical Phase Transitions

*Friday, 21 February 2020 10:00 (1 hour)*

Quantum conductance in molecular and nano systems at low temperatures becomes well understood through the Landauer's motto: "conductance is a transmittance" as calculated from the Schrödinger Equation (SE). However, quantum dynamics is inevitably affected by decoherence which, in turn, becomes critical to recover the macroscopic Ohm's law. In spite of this, the few works that include these effects still resort to the early developments that describe the steady-state in terms of the D'Amato-Pastawski (DP) model [1,2]. Its basic idea is that because of the electron-phonon and other interactions with an "environment", an electron loses its memory of its initial state following a Poisson process with a mean-life described by a Fermi Golden Rule (FGR). After each interaction it initiates a "new life". Here, I will show that non-Hermitian Hamiltonians and the Keldysh Green's functions are the natural setting to describe these processes [3]. I describe how Keldysh integral equations become highly simplified for the DP model in a linear response regime consistent with Landauer approach. The observables are evaluated self-consistently in a discrete space and can be readily generalized to time dependent situations in a multi-terminal settings.[4].

As illustrations for these equations I discuss 1-D model for the Giant Magnetoresistance (GMR). Starting from a quantum description, we recover the standard two resistor model of the GMR. This is achieved by increasing the decoherence controlled by the mean free path and the spin-flip rate, which are Hamiltonian parameters, and determine the observed transport properties. In a device length scale lower than the mean free path, there are interferences that depend on the domain wall size identified with Rabi oscillations. I also discuss a Resonant Tunneling Device where the e-ph scattering processes are considered, within a Fock space description beyond the FGR approximation. This leads to a simple model, which becomes a Floquet Hamiltonian for a classical vibrational field, with antiresonances and resonances in the Fock-Floquet space that describe a phonon laser (SASER) [5]. The inclusion of further decoherence processes, besides smoothing out of the resonances, leads to the degradation of the contrast mainly from the suppression of the antiresonances.

In a second part, I will show that non-Hermitian Hamiltonians not only imply a trivial exponential decay of quantum memory but could also induce a Quantum Dynamical Phase Transitions as that observed in a SWAP gate [6]. These can be interpreted in terms of spectral bifurcations (exceptional points) in the solutions of the Schrödinger equation, i.e. the complex eigen-energies of the non-Hermitian Hamiltonian. However, quite the dynamical phase transition only shows up in the density described by Keldysh or Lindblad equations, as exceptional points in frequency spectrum, not associated with the energies [7]. I also will explore this situation with a stochastic SE algorithm [8], the quantum drift (QD), which is a dynamical implementation of the DP model that captures QDPT. As QD involves a single pure wave function that represents the whole thermal ensemble, it is very effective to push the computational limits for decoherent many-body dynamics. Furthermore, by implementing a forward quantum dynamics followed by another one backwards in time and evaluating the Loschmidt Echo, showing that decoherence does not occur uniformly in time but that it is more effective while the state goes through a superposition (entangled) state. This is consistent with our recent experimental findings [9] that the entangled states are more sensitive to the action of local environments than the simple pointer states. Thus, the extreme sensitivity of the SE of a many-body system to even infinitesimal perturbations, ensures the emergence of the Second Law of Thermodynamics.

[1] J. L. D'Amato and H. M. Pastawski, Conductance of a Disordered Linear Chain including inelastic Scattering events. Phys. Rev. B 41, 7411 (1990).

- [2] S. Datta, A simple kinetic equation for steady-state quantum transport J. Phys.: Condens. Matter 2, 8023 (1990).
- [3] H. M. Pastawski Classical and Quantum Transport from Generalized Landauer-Büttiker Equations II: Time dependent tunneling. Phys. Rev. B 46 4053-4070 (1992).
- [4] C. J. Cattena, L. Fernández-Alcázar, D. Nozaki, R. A. Bustos-Marún, and H. M. Pastawski, Generalized multi-terminal decoherent transport: recursive algorithms and applications to SASER and giant magnetoresistance J. Phys. Condens. Matter 26, 345304 (2014).
- [5] L. E. F. Foa Torres, H. M. Pastawski, and S. S. Makler, Tuning a resonance in the Fock space: Optimization of phonon emission in a resonant tunneling device Phys. Rev. B. 64, 193304 (2001); W. Maryam, A.V. Akimov, R.P. Campion A.J. Kent, Dynamics of a vertical cavity quantum cascade phonon laser structure, Nat. Commun. 4, 2184 (2013).
- [6] G.A. Álvarez, E.P. Danieli, P.R. Levstein, and H.M. Pastawski, Environmentally induced quantum dynamical phase transition in the spin swapping operation J.Chem.Phys. 124, 1 (2006).
- [7] H.M. Pastawski, Revisiting the Fermi Golden Rule: Quantum dynamical phase transition as a paradigm shift, Physica B 398 278 (2007).
- [8] L. J. Fernández-Alcázar and H.M. Pastawski, Decoherent time-dependent transport beyond the Landauer-Büttiker formulation: A quantum-drift alternative to quantum jumps, Phys. Rev. A 91, 022117 (2015).
- [9] C. M. Sánchez, A. K. Chattah, K. X. Wei, L. Buljubasich, P. Cappellaro and H. M. Pastawski, Emergent perturbation independent decay of the Loschmidt Echo in a many-body system. arXiv 1902.06628.

**Session Classification:** SUPER PLENARY TALKS

**Track Classification:** SYMPOSIUM ON SCATTERING, QUANTUM AND CLASSICAL TRANSPORT